arXiv:gr-qc/0102081v1 19 Feb 2001

Hawking Radiation of Dirac Particles in a Variable-mass Kerr Black Hole

Wu Shuang-Qing* and Cai Xu†

Institute of Particle Physics, Hua-Zhong Normal University, Wuhan 430079, China

*E-mail: sqwu@iopp.ccnu.edu.cn †E-mail: xcai@ccnu.edu.cn

(February 7, 2008)

Hawking effect of Dirac particles in a variable-mass Kerr space-time is investigated by using method of the generalized tortoise coordinate transformation. The location and the temperature of event horizon of the non-stationary Kerr black hole are derived. It is shown that the temperature and the shape of event horizon depend not only on the time but also on the polar angle. However, our results demonstrate that the Fermi-Dirac spectrum displays a residual term which is absent from that of Bose-Einstein distribution.

PACS: 04.70.Dy, 97.60.Lf

An important subject in black hole physics is to reveal thermal properties of various black holes. In order to study the Hawking effect [1] of a nonstationary black hole, Zhao and Dai [2] suggested a novel method of the generalized tortoise coordinate transformation (GTCT) which can give simultaneously the exact values both of the location and of the temperature of the event horizon of a non-stationary black hole. The GTCT method has been applied to investigate thermal radiation of scalar particles in the non-stationary Kerr black hole [3] and in the non-stationary Kerr-Newman black hole [4].

However, it is very difficult to discuss on the evaporation of Dirac particles in the non-stationary axisymmetric black hole. The difficulty lies in the non-separability of the radial and angular variables for Chandrasekhar-Dirac equations [5] in the non-stationary axisymmetry spacetime. Though the Hawking effect of Dirac particles in the non-stationary Kerr black hole has been studied in Ref. [6], the location and the temperature of the event horizon they obtained are only valid for the case of slow rotation and small evaporation [7], because their ansatz of separating variables (11) to equation (10) in Ref. [6], strictly speaking, is invalid in the most general case.

In this letter, we try to tackle with the thermal radiation of Dirac particles in the non-stationary Kerr space-time. It is shown that the location and the temperature of event horizon is just the same as that obtained in the case of the thermal radiation of Klein-Gordon scalar field in the non-stationary Kerr space-time, but the Fermionic spectrum of Dirac particles displays another new effect dependent on the interaction between the spin of Dirac particles and the angular momentum of black holes. This has not been reported in the previous studies yet.

The variable mass Kerr solution [8,9] can be written in the advanced Eddington-Finkelstein system as

$$\begin{aligned} ds^2 = &\frac{\Delta - a^2\sin^2\theta}{\Sigma}dv^2 - 2dvdr + 2a\sin^2\theta drd\varphi \\ &- \Sigma d\theta^2 + 2\frac{r^2+a^2-\Delta}{\Sigma}a\sin^2\theta dvd\varphi \\ &- \frac{(r^2+a^2)^2 - \Delta a^2\sin^2\theta}{\Sigma}\sin^2\theta d\varphi^2 \,, \end{aligned} \tag{1}$$

where $\Delta = r^2 - 2M(v)r + a^2$, $\Sigma = \bar{\rho}\rho$, in which $\rho = r - ia\cos\theta$, $\bar{\rho}$ is the complex conjugate of ρ, and v is the standard advanced time. The mass M depends on the time v, but the specific angular momentum a is a constant.

With the choice of a complex null-tetrad such that its corresponding directional derivatives are

$$\begin{aligned} &\underline{D} = -\partial_r \,, \quad \underline{\Delta} = \Sigma^{-1}\Big[(r^2+a^2)\partial_v + 2^{-1}\Delta\partial_r + a\partial_\varphi\Big] \,, \\ &\delta = \frac{1}{\sqrt{2}\bar{\rho}}\left(ia\sin\theta\partial_v + \partial_\theta + \frac{i}{\sin\theta}\partial_\varphi\right) \end{aligned} \tag{2}$$

and $\overline{\delta}$ the complex conjugate of δ, one can obtain the following Dirac equation [5]

$$\begin{aligned} -\left(\partial_r + \tfrac{r}{\Sigma}\right) F_1 + \tfrac{1}{\sqrt{2}\rho}\left(\mathcal{L} - \tfrac{ira\sin\theta}{\Sigma}\right) F_2 &= \tfrac{i\mu_0}{\sqrt{2}} G_1 \,, \\ \tfrac{\Delta}{2\Sigma}\left(\mathcal{D} - \tfrac{ia\cos\theta}{\Sigma}\right) F_2 + \tfrac{1}{\sqrt{2}\bar{\rho}}\left(\bar{\mathcal{L}} - \tfrac{a^2\sin\theta\cos\theta}{\Sigma}\right) F_1 &= \tfrac{i\mu_0}{\sqrt{2}} G_2 \,, \\ -\left(\partial_r + \tfrac{r}{\Sigma}\right) G_2 - \tfrac{1}{\sqrt{2}\bar{\rho}}\left(\bar{\mathcal{L}} + \tfrac{ira\sin\theta}{\Sigma}\right) G_1 &= \tfrac{i\mu_0}{\sqrt{2}} F_2 \,, \\ \tfrac{\Delta}{2\Sigma}\left(\mathcal{D} + \tfrac{ia\cos\theta}{\Sigma}\right) G_1 - \tfrac{1}{\sqrt{2}\rho}\left(\mathcal{L} - \tfrac{a^2\sin\theta\cos\theta}{\Sigma}\right) G_2 &= \tfrac{i\mu_0}{\sqrt{2}} F_1 \,, \end{aligned} \tag{3}$$

where μ_0 is the mass of Dirac particles, the operators have been defined as

$$\begin{aligned} \mathcal{D} &= \partial_r + \Delta^{-1}\left[r - M + 2a\partial_\varphi + 2(r^2 + a^2)\partial_v\right] \,, \\ \mathcal{L} &= \partial_\theta + \frac{1}{2}\cot\theta - \frac{i}{\sin\theta}\partial_\varphi - ia\sin\theta\partial_v \,, \\ \bar{\mathcal{L}} &= \partial_\theta + \frac{1}{2}\cot\theta + \frac{i}{\sin\theta}\partial_\varphi + ia\sin\theta\partial_v \,. \end{aligned}$$

By substituting

$$F_1 = \frac{P_1}{\sqrt{2\Sigma}} \,, \; F_2 = \frac{\rho P_2}{\sqrt{\Sigma}} \,, \; G_1 = \frac{\bar{\rho} Q_1}{\sqrt{\Sigma}} \,, \; G_2 = \frac{Q_2}{\sqrt{2\Sigma}} \,,$$

into Eq. (3), they have the form

$$\begin{aligned} &-\partial_r P_1 + \mathcal{L} P_2 = i\mu_0\bar{\rho} Q_1 \,, \;\; \Delta\mathcal{D} P_2 + \bar{\mathcal{L}} P_1 = i\mu_0\bar{\rho} Q_2 \,, \\ &-\partial_r Q_2 - \bar{\mathcal{L}} Q_1 = i\mu_0\rho P_2 \,, \;\; \Delta\mathcal{D} Q_1 - \mathcal{L} Q_2 = i\mu_0\rho P_1 \,. \end{aligned} \tag{4}$$

Eq. (4) can not be decoupled except in the case of Kerr black hole ($M = \text{const}$) [5]. However, to deal with the problem of Hawking radiation, one may concern about the behavior of Eq. (4) near the horizon only. Now, let us make a GTCT [2]

$$\begin{aligned} r_* &= r + \tfrac{1}{2\kappa_H}\ln(r - r_H) \,, \\ v_* &= v - v_0 \,, \;\; \theta_* = \theta - \theta_0 \,, \end{aligned} \tag{5}$$

where $r_H = r_H(v, \theta)$ is the location of the event horizon, and κ_H is an adjustable parameter. All parameters κ_H, v_0 and θ_0 are constant under the tortoise transformation.

Under the transformations (5), Eq. (4) can be reduced to

$$\begin{aligned} &\partial_{r_*} P_1 + (r_H' - ia\sin\theta_0\dot{r}_H)\,\partial_{r_*} P_2 = 0 \,, \\ &[\Delta_H - 2(r_H^2 + a^2)\dot{r}_H]\,\partial_{r_*} P_2 \\ &\qquad - (r_H' + ia\sin\theta_0\dot{r}_H)\,\partial_{r_*} P_1 = 0 \,, \end{aligned} \tag{6}$$

after being taken limits $r \to r_H(v_0, \theta_0)$, $v \to v_0$ and $\theta \to \theta_0$. In the above, we have denoted $\dot{r}_H = \partial_v r_H$, $r_H' = \partial_\theta r_H$, and $\Delta_H = r_H^2 - 2M(v_0) r_H + a^2$.

If the derivatives $\partial_{r_*}P_1$ and $\partial_{r_*}P_2$ in Eq. (6) are not equal to zero, the existence condition of non-trial solutions for P_1 and P_2 is that the determinant of Eq. (6) vanishes, which gives the following equation to determine the location of horizon

$$\Delta_H - 2(r_H^2 + a^2)\dot{r}_H + a^2\sin^2\theta_0\dot{r}_H^2 + {r'_H}^2 = 0\,. \tag{7}$$

An apparent fact is that the Chandrasekhar-Dirac equations (4) could be satisfied by identifying Q_1, Q_2 with $\bar{P}_2$, $-\bar{P}_1$, respectively. So one may deal with a pair of components P_1, P_2 only. Now let us consider the asymptotic behaviors of the second-order equations for the two-component spinor (P_1, P_2) near the event horizon. Given the GTCT in Eq. (5), the limiting form of the second order equations for P_1, P_2, when r approaches $r_H(v_0, \theta_0)$, v goes to v_0 and θ goes to θ_0, reads

$$\begin{aligned}
&\mathcal{K}P_1 - \{r_H(1 - 3\dot{r}_H) - M + r''_H + \ddot{r}_H a^2\sin^2\theta_0\\
&+\ r'_H\cot\theta_0 + \frac{\Delta_H - \dot{r}_H(r_H^2 + a^2) + ia\sin\theta_0 r'_H}{\bar{\rho}_H}\}\partial_{r_*}P_1\\
&+\ \{2i\dot{M}r_H a\sin\theta_0 - \frac{2ia\sin\theta_0\dot{r}_H(r_H^2 + a^2)}{\bar{\rho}_H}\\
&-\ \frac{\Delta_H[r'_H - ia\sin\theta_0(\dot{r}_H + 1)]}{\bar{\rho}_H}\}\partial_{r_*}P_2 = 0\,,
\end{aligned} \tag{8}$$

and

$$\begin{aligned}
&\mathcal{K}P_2 - \{M - r_H(1 + \dot{r}_H) + r''_H + \ddot{r}_H a^2\sin^2\theta_0\\
&+\ r'_H\cot\theta_0 + \frac{\Delta_H - \dot{r}_H(r_H^2 + a^2) + ia\sin\theta_0 r'_H}{\bar{\rho}_H}\}\partial_{r_*}P_2\\
&+\ \frac{r'_H + ia\sin\theta_0(\dot{r}_H - 1)}{\bar{\rho}_H}\partial_{r_*}P_1 = 0\,.
\end{aligned} \tag{9}$$

where $\bar{\rho}_H = r_H + ia\cos\theta_0$. We have used the L' Hôsptial rule to get a term involving the second-order derivatives denoted by the operator

$$\begin{aligned}
\mathcal{K} = &\Big\{\kappa_H^{-1}[r_H(1 - 2\dot{r}_H) - M] - 2\dot{r}_H(r_H^2 + a^2)\\
&+\ 2\Delta_H\Big\}\,\partial_{r_*}^2 + 2(r_H^2 + a^2 - \dot{r}_H a^2\sin^2\theta_0)\partial_{r_*v_*}^2\\
&+\ 2a(1 - \dot{r}_H)\partial_{r_*\varphi}^2 P_2 - 2r'_H\partial_{r_*\theta_*}^2
\end{aligned}$$

One can adjust the parameter κ_H such that it satisfies

$$\begin{aligned}
&\kappa_H^{-1}[r_H(1 - 2\dot{r}_H) - M] + 2\Delta_H - 2\dot{r}_H(r_H^2 + a^2)\\
&\equiv\ r_H^2 + a^2 - \dot{r}_H a^2\sin^2\theta_0\,,
\end{aligned} \tag{10}$$

which means the temperature of the horizon is

$$\kappa_H = \frac{r_H(1 - 2\dot{r}_H) - M}{r_H^2 + a^2 - \dot{r}_H a^2\sin^2\theta_0 - 2\Delta_H + 2\dot{r}_H(r_H^2 + a^2)}\,. \tag{11}$$

Using Eq. (6), Eq. (8) and Eq. (9) can be recast into the standard wave equation near the horizon in an united form

$$\begin{aligned} &\partial^2_{r_*}\Psi + 2\partial^2_{r_* v_*}\Psi + 2\Omega\partial^2_{r_*\varphi}\Psi + 2C_3\partial^2_{r_*\theta_*}\Psi \\ &\quad + \ 2(C_2 + iC_1)\partial_{r_*}\Psi = 0\,, \end{aligned} \tag{12}$$

where $\Omega = a(1 - \dot{r}_H)/(r_H^2 + a^2 - \dot{r}_H a^2 \sin^2\theta_0)$ is the angular velocity of the event horizon of the evaporating Kerr black hole. $C_3 = -r_H'/(r_H^2 + a^2 - \dot{r}_H a^2 \sin^2\theta_0)$, C_2 (omitted here) and C_1 are all real constants,

$$\begin{aligned} C_1 &= \frac{1}{2(r_H^2 + a^2 - \dot{r}_H a^2 \sin^2\theta_0)}[-\dot{r}_H a\cos\theta_0 \\ &\quad + \frac{2\dot{M} r_H r_H' a \sin\theta_0}{\Delta_H - 2\dot{r}_H(r_H^2 + a^2)}]\,, \qquad \text{for} \quad \Psi = P_1, \qquad (13) \\ C_1 &= \frac{\dot{r}_H a\cos\theta_0}{2(r_H^2 + a^2 - \dot{r}_H a^2 \sin^2\theta_0)}\,, \qquad \text{for} \quad \Psi = P_2. \qquad (14) \end{aligned}$$

Now separating as $\Psi = R(r_*)\exp[\lambda\theta_* + i(m\varphi - \omega v_*)]$, one has

$$\begin{aligned} &R'' + 2(C_0 + iC_1 + im\Omega - i\omega)R' = 0\,, \\ &R \sim e^{2i(\omega - m\Omega - C_1)r_* - 2C_0 r_*}\,; \quad R_0\,. \end{aligned} \tag{15}$$

where $C_0 = \lambda C_3 + C_2$, λ is a real constant introduced in the separation of variables.

The ingoing wave solution and the outgoing wave solution to Eq. (12), are respectively,

$$\begin{aligned} &\Psi_{\rm in} = \exp[-i\omega v_* + im\varphi + \lambda\theta_*]\,, \\ &\Psi_{\rm out} = \Psi_{\rm in} e^{2i(\omega - m\Omega - C_1)r_* - 2C_0 r_*}\,, \quad (r > r_H)\,. \end{aligned} \tag{16}$$

The outgoing wave $\Psi_{\rm out}$ is not analytic at the event horizon $r = r_H$, but can be analytically continued from the outside of the hole into the inside of the hole by the lower complex r-plane to

$$\tilde{\Psi}_{\rm out} = \Psi_{\rm out} e^{i\pi C_0/\kappa_H} e^{\pi(\omega - m\Omega - C_1)/\kappa_H} \quad (r < r_H)\,. \tag{17}$$

The relative scattering probability at the event horizon is

$$\left|\frac{\Psi_{\rm out}}{\tilde{\Psi}_{\rm out}}\right|^2 = e^{-2\pi(\omega - m\Omega - C_1)/\kappa_H}\,. \tag{18}$$

Following the method of Damour-Ruffini-Sannan's [10], the Fermionic spectrum of Hawking radiation of Dirac particles from the black hole is easily obtained

$$\langle\mathcal{N}(\omega)\rangle = \frac{1}{e^{(\omega - m\Omega - C_1)/T_H} + 1}\,, \quad T_H = \frac{\kappa_H}{2\pi}\,. \tag{19}$$

Equations (7) and (11) give the location and the temperature of event horizon of the variable-mass Kerr black hole, which depend not only on the advanced time v but also on the polar angle θ. They are in accord with that obtained from Klein-Gordon field equation [3,4]. Eq. (19) shows the thermal radiant spectrum of Dirac particles in the non-stationary Kerr space-time, in which a residual term C_1 appears. The difference between Bosonic spectrum and Fermionic spectrum is that C_1 is absent in the spectrum of Klein-Gordon scalar particles. Also C_1 vanishes when a black hole is stationary or has a vanishing angular momentum.

In conclusion, the C_1 term represents a new effect in the spectrum of Dirac particles, which is absent from the spectrum of Klein-Gordon particles. This new effect maybe arise from the interaction between the spin of Dirac particle and the evaporating black hole.

S.Q. Wu is very grateful to Dr. Jeff Zhao at Motomola Company for his longterm helps. This work is supported in part by the NSFC under Grant No.19875019.

---

[1] Hawking S W 1974 *Nature* **248** 30; 1975 *Commun. Math. Phys.* **43** 199

[2] Zhao Z and Dai X X 1991 *Chin. Phys. Lett.* **8** 548

[3] Zhao Z, Dai X X and Huang W H 1993 *Acta Astrophysica Sinica* **13** 299

[4] Luo M W 2000 *Acta Physica Sinica* **49** 1035; Jing J L and Wang Y J 1997 *Int. J. Theor. Phys.* **36** 1745

[5] Chandrasekhar S 1983 *The Mathematical Theory of Black Holes* (New York: Oxford University Press)

[6] Li Z H and Zhao Z 1995 *Science in China* A **38** 74

[7] Zhao Z and Huang W H 1992 *Chin. Phys. Lett.* **9** 333

[8] Gonzalez C, Herrera L and Jimenez J 1979 *J. Math. Phys.* **20** 837; Jing J L and Wang Y J 1996 *Int. J. Theor. Phys.* **35** 1841

[9] Carmeli M and Kaye M 1977 *Ann. Phys.* **103** 197; Carmeli M 1982 *Classical Fields: General Relativity and Gauge Theory* (New York: John Wiley Sons)

[10] Damour T and Ruffini R 1976 *Phys. Rev.* D **14** 332; Sannan S 1988 *Gen. Rel. Grav.* **20** 239